

## Genetic Monogamy in Carolina Wrens (*Thryothorus ludovicianus*)

THOMAS M. HAGGERTY,<sup>1,4</sup> EUGENE S. MORTON,<sup>2</sup> AND ROBERT C. FLEISCHER<sup>3</sup>

<sup>1</sup>*Department of Biology, University of North Alabama, Florence, Alabama 35632, USA;*

<sup>2</sup>*Conservation and Research Center, National Zoological Park, 1500 Remount Road,  
Front Royal, Virginia 22630, USA; and*

<sup>3</sup>*National Zoological Park, Smithsonian Institution, Washington, D.C. 20008, USA*

Molecular comparisons have shown that socially monogamous passerines often have mixed reproductive strategies (Birkhead and Møller 1992, 1996). Pairs often cooperate in raising a brood, but each sex may pursue additional extrapair matings (e.g. Westneat 1987, Morton et al. 1990, Kempenaers et al. 1992). Further, females of some species lay eggs in nests of conspecifics (i.e. intraspecific brood parasitism, ISBP; reviewed in Hughes 1998).

Although considerable intra- and interspecific variation has been found in rates of extrapair paternity

(EPP), causes for that variation remain unclear and additional data are warranted (Petrie and Kempenaers 1998). Further, few studies have been conducted on temperate-zone species that defend a territory and maintain a pair bond year round. In this study, we use multilocus DNA fingerprinting to examine paternity and intraspecific brood parasitism in Carolina Wrens (*Thryothorus ludovicianus*), a socially monogamous species that maintains a year-round pair bond and territory (Haggerty and Morton 1995). In addition, we report on breeding synchrony in Carolina Wrens because it is an ecological factor that may be related to paternity (Møller and Ninni 1998).

---

<sup>4</sup> E-mail: thaggert@unanov.una.edu

**Methods.**—The study was conducted between March and August, 1996 and 1997, on a 43 ha mixed hardwood forest on the Tennessee Valley Authority reservation in Muscle Shoals, Colbert County, Alabama (34°49'N, 87°38'W). The overstory and understorey are dominated by hackberry (*Celtis laevigata*) and privet (*Ligustrum vulgare*), respectively. During most of the breeding season, the ground-cover vegetation is dominated by honeysuckle (*Lonicera japonica*), poison ivy (*Rhus radicans*), and Virginia creeper (*Parthenocissus quinquefolia*).

Nest boxes were provided in late winter (5–6 per territory) and were readily used by Carolina Wrens (Haggerty and Morton 1995). Adults were captured near their nests with mist nets and approximately 30 to 100  $\mu$ L of blood was collected from the brachial vein, stored in phosphate buffered saline/EDTA buffer (1996) or a lysis buffer (1997) and refrigerated. Blood samples from nestlings were collected in a similar way when they were 5 to 8 days old (hatching day = day 0). Adults were marked with a unique combination of colored leg bands and a U.S. Fish and Wildlife aluminum band. Parents caring for nestlings were the putative parents. Most adults had been previously banded as part of a long-term population study that began in 1988 (Haggerty and Morton 1995). Age (i.e. number of breeding years on study area) for adults that were fingerprinted ranged from 1 to 5 years ( $\bar{x} = 1.6$ ).

A 50  $\times$  50 m grid system was established to help calculate size of the study area and to determine pair density. The 1996 and 1997 breeding-pair densities were 4.2 individuals/10 ha and 7.9/10 ha, respectively. A breeding-synchrony index was calculated for each year (Kempnaers 1993).

Multilocus DNA fingerprinting was conducted following the protocol of Loew and Fleischer (1996) using the Jeffreys 33.15 probe (Jeffreys et al. 1985). *Hae*III digested DNA of nestlings was usually placed in lanes between their putative parents for ease of scoring. Resulting autoradiographs were scored by counting number of fragments in a nestling's lane that were attributable to either or both parents profiles, and the number that were not (i.e. novel fragments). Pairwise band-sharing coefficients (*S*) were calculated according to Lynch (1991). A total of 84 offspring and 32 putative parents from 23 nests (i.e. a total of 116 individuals) were fingerprinted. Seven pairs were scored for two nests and nine pairs for only one nest. All offspring were fingerprinted for 16 of the 23 nests, but eight nestlings in seven nests were not analyzed because DNA was degraded or some other technical problem. Mean number of fragments per individual profile was  $13.3 \pm$  SD of 3.8 (range 6 to 24). Only fragments above about 3 kb were scored; hence, the smaller than normal number of fragments per profile. Typically, fragments below 3 kb were less variable than larger ones and they




FIG. 1. Frequency of novel fragments among Carolina Wren offspring compared to their putative parents. Bars represent the observed frequencies. The line represents the theoretical distribution calculated from a Poisson distribution on the basis of mean number of fragments (0.238,  $n = 84$ ) from nestlings with fewer than four novel fragments.

added little or no information to estimates of relatedness.

**Results.**—The synchrony indices for 1996 and 1997 were  $17.7\% \pm 10$  ( $n = 11$  nests, 5 females) and  $14.3\% \pm 11.0$  ( $n = 39$  nests, 19 females), respectively.

All DNA fragments found in offspring profiles were also found in the parent's profiles for 68 of the 84 offspring. For 13 offspring, we found one novel or nonattributable fragment. For two offspring, we found two novel bands and for one we found three. Mean number of novel fragments was  $0.238 \pm 0.55$ , corresponding to a mutation or artifact rate of 0.019 per fragment/generation. The distribution of novel fragments matches a Poisson expectation, on the basis of a mean of 0.238 ( $n = 84$  profiles; Fig. 1). Because that match suggests that mutation alone can explain the extra fragments, we concluded, following the rationale of Westneat (1990), that there were no extra-pair fertilizations (EPFs) in our sample.

The mean value of *S* calculated for the 16 pairs of parents was  $0.225 \pm 0.13$ , which did not differ significantly from 11 random pairwise *S* values ( $\bar{x} = 0.24 \pm 0.1$ ;  $t = 0.23$ ,  $P = 0.82$  [or Mann-Whitney  $U = 82.0$ ,  $P = 0.77$ ]). The predicted mean *S* for first-order relatives ( $R = 0.5$ ) was 0.59 (equation 22 of Lynch 1991). Based on the level of background band-sharing, the probability of assigning parents incorrectly (ISBP) was  $5.9 \times 10^{-6}$ , whereas the probability of assigning the male parent incorrectly (EPF) was  $1.9 \times 10^{-4}$  (Bruford et al. 1998). The mean value of *S* for 84 comparisons of female parents and offspring was  $0.52 \pm 0.13$ , whereas *S* for male parents and offspring was  $0.55 \pm 0.11$  (Fig. 2). The plot of *S* against




FIG. 2. Band-sharing among putative Carolina Wren parents and nestlings. Symbol location denotes the proportion of bands in a nestling's fingerprint shared with putative mother and father, and plotted against the number of bands in the nestling's fingerprint that were not in the putative parents' fingerprint (novel fragments).

number of novel fragments shows that those individuals with 1, 2, or 3 novel fragments have high levels of band-sharing (Fig. 2), providing additional evidence that the 84 nestlings in 23 nests cannot be excluded from the adults attending those nests.

**Discussion.**— We found no evidence of a mixed reproductive strategy in our population of Carolina

Wrens. The lack of ISBP was expected because we did not find any nests in which more than one egg had been laid over a 24 h period.

Factors that may affect opportunities for EPFs include breeding synchrony (Stutchbury and Neudorft 1998), population density (Westneat et al. 1990, Westneat and Sherman 1997, Møller and Ninni 1998), and mate guarding (Westneat et al. 1990, Currie et al. 1999). Westneat (1990) proposed that breeding synchrony should reduce frequency of EPP because males would be too busy guarding mates to engage in extrapair copulations (EPCs). Stutchbury and Morton (1995), however, proposed that breeding synchrony allows females to evaluate male quality and promotes EPP in some species. Our population had an overall low synchrony index value (i.e.  $15.4\% \pm 11$ ), which supports the Stutchbury and Morton (1995) hypothesis.

Although population densities during the years of this research were not the highest observed on the study site (i.e. 15 individuals/10 ha), territorial boundaries expand and are often shared even during low-density years (T. Haggerty pers. obs.). Therefore, we suspect that opportunities for EPFs existed and that a low density was not the primary cause for genetic monogamy in our population.

Although mate guarding may have occurred in our population, fledgling care should have limited the male's ability to guard their mates during the fertile periods of subsequent broods (Weatherhead and McRae 1990; but see Møller 1991, Conrad et al. 1998). Yet, we found no extrapair young in subsequent broods (i.e. 7 nests, 27 nestlings). Furthermore, territorial advertisement and defense in a visually occluded habitat should have made mate guarding difficult (Westneat et al. 1990) and some EPFs possible, yet none was recorded. Our population also has a low divorce rate (i.e. 2 of 36 cases from 27 pairs over 12 breeding seasons in which both pair members survived from one breeding season to the next) which is contrary to what is predicted when monogamy is enforced (Birkhead and Møller 1996, Gowaty 1996), but is expected when EPP rates are low (Cezilly and Nager 1995). Therefore, we doubt that mate guarding constrained females from engaging in extrapair activities.

As expected for a species with a low EPF rate (Birkhead and Møller 1996), males in our population contributed substantially to offspring care (Haggerty and Morton 1995, T. Haggerty unpubl. data). Although males do not incubate, they provide food to nestlings and females. In addition, nesting-interval data (Haggerty and Morton 1995) show that females lay and incubate new clutches well before fledglings from previous broods reach independence. Therefore, male care during the fledgling period may be essential if multiple broods are to be raised by a pair during a prolonged breeding season (Westneat et al. 1990). The threat of male desertion or reduced care

may constrain females from engaging in EPCs (Møller 1988, Burke et al. 1989, Dixon et al. 1994). Further, in sedentary species like the Carolina Wren, females may need mutually defended resources year round for their survival. Females engaging in EPCs may lose access to defended resources (Møller 1988, Westneat and Gray 1998). Most Carolina Wren mortality occurs during winter (Haggerty and Morton 1995, T. Haggerty unpubl. data) and females that have a mate may have a better chance of surviving and breeding another year than unfaithful and unmated females.

*Acknowledgments.*—We thank S. Hardin and the many other students who have assisted T. Haggerty in his field work over the years. Thanks to B. Stutchbury and M. Moeller for their assistance in the calculation of the synchrony indices. This research was supported by faculty research grants from the University of North Alabama to T. Haggerty and by a Scholarly Studies Grant from the Smithsonian Institution to E. Morton.

## LITERATURE CITED

- BIRKHEAD, T. R., AND A. P. MØLLER. 1992. Sperm Competition in Birds: Evolutionary Causes and Consequences. Academic Press, London.
- BIRKHEAD, T. R., AND A. P. MØLLER. 1996. Monogamy and sperm competition in birds. Pages 323–343 in *Partnerships in Birds* (J. M. Black, Ed.). Oxford University Press, New York.
- BRUFORD, M. W., O. HANNOTTE, AND T. BURKE. 1998. Multi-locus and single-locus DNA fingerprinting. Pages 287–336 in *Molecular Genetic Analysis of Populations* (A. R. Hoelzel, Ed.). IRL Press, Oxford.
- BURKE, T., N. B. DAVIES, M. W. BRUFORD, AND B. J. HATCHWELL. 1989. Parental care and mating behaviour of polyandrous Dunnocks *Prunella modularis* related to paternity by DNA fingerprinting. *Nature* 338:249–251.
- CEZILLY, F., AND R. G. NAGER. 1995. Comparative evidence for a positive association between divorce and extra-pair paternity in birds. *Proceedings of the Royal Society of London Series B* 262:7–12.
- CONRAD, K. F., R. J. ROBERTSON, AND P. T. BOAG. 1998. Frequency of extrapair young increases in second broods of Eastern Phoebes. *Auk* 115:497–502.
- CURRIE, D., A. P. KRUPA, T. BURKE, AND D. B. THOMPSON. 1999. The effect of experimental male removals on extrapair paternity in the Wheatear, *Oenanthe oenanthe*. *Animal Behaviour* 57:145–152.
- DIXON, A., D. ROSS, S. L. C. O'MALLEY, AND T. BURKE. 1994. Paternal investment inversely related to degree of extra-pair paternity in the Reed Bunting. *Nature* 371:698–700.
- GOWATY, P. A. 1996. Battles of the sexes and origins of monogamy. Pages 21–52 in *Partnerships in Birds* (J. M. Black, Ed.). Oxford University Press, New York.
- HAGGERTY, T. M., AND E. S. MORTON. 1995. The Carolina Wren (*Troglodytes ludovicianus*). In *The Birds of North America*, no. 188 (A. Poole and F. Gill, Eds.). Academy of Natural Sciences, Philadelphia, and American Ornithologists' Union, Washington, D.C.
- HUGHES, C. 1998. Integrating molecular techniques with field methods in studies of social behavior: A revolution results. *Ecology* 79:383–399.
- JEFFREYS, A. J., V. WILSON, AND S. L. THEIN. 1985. Hypervariable "minisatellite" regions in human DNA. *Nature* 314:67–73.
- KEMPENAERS, B. 1993. The use of a breeding synchrony index. *Ornis Scandinavica* 24:84.
- KEMPENAERS, B., G. R. VERHEYEN, M. VAN DEN BROECK, T. BURKE, AND A. A. DHONDY. 1992. Extra-pair paternity results from female preference for high-quality males in the Blue Tit. *Nature* 357:494–496.
- LOEW, S., AND R. C. FLEISCHER. 1996. Multilocus DNA fingerprinting. Pages 456–461 in *Molecular Zoology: Advances, Strategies and Protocols* (J. D. Ferraris and S. R. Palumbi, Eds.). Wiley-Liss, New York.
- LYNCH, M. 1991. The similarity index and DNA fingerprinting. *Molecular Biology and Evolution* 7: 478–484.
- MØLLER, A. P. 1988. Paternity and parental care in the swallow, *Hirundo rustica*. *Animal Behaviour* 36: 996–1005.
- MØLLER, A. P. 1991. Double broodedness and mixed reproductive strategies by female swallows. *Animal Behaviour* 42:671–679.
- MØLLER, A. P., AND P. NINNI. 1998. Sperm competition and sexual selection: A meta-analysis of paternity studies of birds. *Behavioral Ecology and Sociobiology* 43:345–358.
- MORTON, E. S., L. FORMAN, AND N. BRAUN. 1990. Extrapair fertilizations and the evolution of colonial breeding in Purple Martins. *Auk* 107:275–283.
- PETRIE, M., AND B. KEMPENAERS. 1998. Extra-pair paternity in birds: Explaining variation between species and populations. *Trends in Ecology and Evolution* 13:52–58.
- STUTCHBURY, B. J., AND E. S. MORTON. 1995. The effect of breeding synchrony on extra-pair mating systems in songbirds. *Behaviour* 132:675–690.
- STUTCHBURY, B. J., AND D. L. NEUDORF. 1998. Female control, breeding synchrony, and the evolution of extra-pair mating tactics. Pages 103–121 in *Male and Female Extra-pair Mating Strategies in Birds* (P. Parker and N. Burley, Eds.). *Ornithological Monographs* no. 49.

- WEATHERHEAD, P. J., AND S. B. MCRAE. 1990. Brood care in American Robins: Implications for mixed reproductive strategies by females. *Animal Behaviour* 39:1179–1188.
- WESTNEAT, D. F. 1987. Extrapair fertilizations in a predominantly monogamous bird: Genetic evidence. *Animal Behaviour* 35:877–886.
- WESTNEAT, D. F. 1990. Genetic parentage in the Indigo Bunting: A study using DNA fingerprinting. *Behavioral Ecology and Sociobiology* 27:67–76.
- WESTNEAT, D. F., AND E. M. GRAY. 1998. Breeding synchrony and extrapair fertilizations in two populations of Red-winged Blackbirds. *Behavioral Ecology* 9:456–464.
- WESTNEAT, D. F., AND P. W. SHERMAN. 1997. Density and extra-pair fertilizations in birds: A comparative analysis. *Behavioral Ecology and Sociobiology* 41:205–215.
- WESTNEAT, D. F., P. W. SHERMAN, AND M. L. MORTON. 1990. The ecology and evolution of extra-pair copulations in birds. *Current Ornithology* 7: 331–369.

*Received 11 October 1999, accepted 5 September 2000.*

*Associate Editor: E. Sheldon*